

1

2

3

4

5

# **Pre-Hospital Sepsis Screening Tool**

UNDER 5 YEARS	AGE 5-11 YEARS	AGE 12+ YEARS (ADULT)	PREGNANT (OR UP TO 6 WE
<ul> <li>START THIS CHART IF PATIENT LOOKS UNWELL, PARENTAL CONCERN, EWS ≥ 5 OR ABNORMAL TEMPERATURE (&lt;36°C OR &gt;38°C)</li> <li>Sepsis risk factors <ul> <li>Recent trauma/surgery/invasive procedure</li> <li>Impaired immunity (e.g. diabetes, steroids, chemotherapy, neutropenic patient, asplenia)</li> <li>Indwelling lines/broken skin/indwelling devices</li> <li>Rural, remote or low socioeconomic status</li> <li>Re-presentation or delayed presentation to a health service</li> <li>Infants &lt; 3 months old</li> </ul> </li> </ul>	<ul> <li>START THIS CHART IF PATIENT LOOKS UNWELL, PARENTAL CONCERN, EWS ≥ 5 OR ABNORMAL TEMPERATURE (&lt;36°C OR &gt;38°C)</li> <li>Sepsis risk factors <ul> <li>Recent trauma/surgery/invasive procedure</li> <li>Impaired immunity (e.g. diabetes, steroids, chemotherapy, neutropenic patient, asplenia)</li> <li>Indwelling lines/broken skin/indwelling devices</li> <li>Rural, remote or low socioeconomic status</li> <li>Re-presentation or delayed presentation to a health service</li> </ul> </li> </ul>	<ul> <li>START THIS CHART IF PATIENT LOOKS UNWELL OR EWS ≥ 5</li> <li>Sepsis risk factors <ul> <li>Recent trauma/surgery/invasive procedure</li> <li>Impaired immunity (e.g. diabetes, steroids, chemotherapy, neutropenic patient)</li> <li>Indwelling lines/IVDU/broken skin/indwelling devices</li> <li>Rural, remote or low socioeconomic status</li> <li>AGE &gt;75</li> </ul> </li> </ul>	<ul> <li>START THIS CHART IF PATIENT L OR EWS ≥ 5</li> <li>Sepsis risk factors <ul> <li>Recent trauma/surgery/invasive</li> <li>Impaired immunity (e.g. diabete chemotherapy, neutropenic pat</li> <li>Indwelling lines/IVDU/broken sidevices</li> <li>Rural, remote or low socioecond</li> </ul> </li> </ul>
ANY OBVI	OUS NON-INFECTIVE CAUSE FOR THE PATIEN	NT'S PRESENTATION AND OBSERVATIONS? N	O↓YES→
Sepsis red flags include:	Sepsis red flags include:	Sepsis red flags include:	Sepsis red flags include:
<ul> <li>BSL ≤ 3 mmol/L (note: &lt;2.6 mmol/L for neonates)</li> <li>Doesn't wake when roused or won't stay awake</li> <li>Child looks unwell to healthcare professional</li> <li>Weak, high pitched or continuous cry</li> <li>Tachypnoea (see chart)</li> <li>Tachycardia (see chart)</li> <li>Bradycardia (see chart)</li> <li>Hypotension (see chart)</li> <li>SpO2 ≤ 88% on air or O2 requirements &gt;10 litres/ minute</li> <li>Non-blanching rash</li> <li>Lactate &gt;4mmol/L (if available)</li> </ul>	<ul> <li>BSL ≤3mmol/L</li> <li>Doesn't wake when roused or won't stay awake</li> <li>Objective evidence of new or altered mental state</li> <li>Child looks unwell to healthcare professional</li> <li>Tachypnoea ≥50 breaths/minute</li> <li>Tachycardia ≥170 beats/minute</li> <li>Bradycardia &lt;60 beats/minute</li> <li>Hypotension (systolic BP &lt;60 mmHg)</li> <li>SpO2 ≤ 88% on air or O2 requirements ≥ 10 litres/ minute</li> <li>Non-blanching rash</li> <li>Lactate &gt;4mmol/L (if available)</li> </ul>	<ul> <li>Objective evidence of new or altered mental state</li> <li>Systolic BP ≤90mmHg (or &gt;40mmHg drop from normal)</li> <li>Heart rate ≥130 per minute</li> <li>Temperature ≤ 35°C</li> <li>Respiratory rate ≥25 per minute</li> <li>Needs O2 to keep SpO2 ≥94% (88% in COPD)</li> <li>Non-blanching rash/mottled/ashen/cyanotic</li> <li>Not passed urine in 18 hrs (0.5mL/kg/hr if catheterised)</li> <li>Lactate ≥2 (if available)</li> <li>Recent chemotherapy</li> </ul>	<ul> <li>Objective evidence of new or alt</li> <li>Systolic BP ≤90mmHg (or &gt;40mmHg drop from norma</li> <li>Heart rate ≥130 per minute</li> <li>Temperature ≤ 35°C</li> <li>Respiratory rate ≥25 per minute</li> <li>Needs O2 to keep SpO2 ≥94%</li> <li>Non-blanching rash/mottled/as</li> <li>Not passed urine in 18 hrs (&lt;0.5 catheterised)</li> <li>Lactate ≥2 mmol/L (if available) be raised in and immediately after the second secon</li></ul>
	ANY RED FLAGS PR	ESENT? NO ↓ YES →	
<ul> <li>Sepsis amber flags include:</li> <li>Family and/or clinician concern is continuing or increasing</li> <li>Not responding normally/no smile/reduced activity/very sleepy</li> <li>Unexplained pain, leg pain or cold extremities</li> <li>Tachypnoea (see chart)</li> <li>Tachycardia (see chart)</li> <li>Hypotension (see chart)</li> <li>SpO2 ≤ 91% on air or O2 requirements ≥5 litres/minute</li> <li>Capillary refill time ≥ 3 seconds, mottled or cold extremities</li> <li>Lactate 2-4 mmol/L (if available)</li> <li>Nasal flaring</li> <li>Reduced urine output</li> </ul>	<ul> <li>Sepsis amber flags include:</li> <li>Family and/or clinician concern is continuing or increasing</li> <li>Behaving abnormally/not wanting to play/drowsy/confused</li> <li>Unexplained pain, leg pain or cold extremities</li> <li>Tachypnoea 45-49 breaths per minute</li> <li>Tachycardia 160-169 beats per minute</li> <li>Hypotension (systolic BP &lt;70mmHg)</li> <li>SpO2 ≤ 91% on air or O2 requirements ≥5 litres/minute</li> <li>Capillary refill time ≥ 3 seconds, mottled or cold extremities</li> <li>Lactate 2-4 mmol/L (if available)</li> <li>Reduced urine output (&lt;1mL/kg/hr if catheterised)</li> </ul>	<ul> <li>Sepsis amber flags include:</li> <li>Relatives concerned about mental status</li> <li>Acute deterioration in functional ability</li> <li>Trauma/surgery/procedure in past 8 weeks</li> <li>Immunosuppressed</li> <li>Respiratory rate 21-24 per minute</li> <li>Systolic BP 91-100mmHg</li> <li>Heart rate 91-129 per minute or new dysrhythmia</li> <li>Temperature range between 35.1°C- 36°C or ≥ 38.1°C</li> <li>Clinical signs of wound infection</li> </ul>	<ul> <li>Sepsis amber flags include:</li> <li>If immunity impaired treat as red f</li> <li>Behavioural/mental status char</li> <li>Acute deterioration in functional</li> <li>Respiratory rate 21-24 per minut</li> <li>Systolic BP 91-100mmHg</li> <li>Heart rate 91-129 per minute or</li> <li>Had an invasive procedure in th weeks (e.g. Cervical cerclage, for miscarriage, termination)</li> <li>Temperature range between 35 ≥38.1°C</li> <li>Has diabetes or gestational dia</li> <li>Close contact with GAS (Group</li> <li>Prolonged rupture of membrand</li> <li>Bleeding/wound infection</li> <li>Offensive vaginal discharge</li> <li>Non-reassuring CTG/foetal tace</li> </ul>
Tachypnoea         Tachycardia         Bradycardia         Hypotension           <3 mths			
<3 mths ≥75 65-74 ≥180 170-179 ≤90 ≤100 ≤50 ≤60	ANY AMBER FLAGS P	RESENT? NO ↓ YES →	

## VEEKS PP)

#### LOOKS UNWELL

sive procedure etes, steroids, patient) n skin/indwelling

nomic status



#### **SEPSIS UNLIKELY**

altered mental state

nal)

ite % (88% in COPD) ashen/cyanotic ).5mL/kg/hr if

e) note: lactate may Ifter normal delivery

# **RED FLAG SEPSIS**

1. Pre-notify ED

- 2. Follow RED EWS escalation advice
- 3. Communicate RED flag sepsis at handover

#### flag sepsis

ange nal ability nute

or new dysrythmia the past 6 forceps delivery,

35.1°C- 36°C or

iabetes up A Streptococcus) ines

achycardia >160



## **AMBER FLAG SEPSIS**

- . Follow AMBER EWS escalation advice
- 2. Communicate AMBER flag sepsis at handover

iagnosis.

Uncontrolled once printed |Oct 2023